

AD-A195 802

④

CHEMICAL
RESEARCH,
-DEVELOPMENT &
ENGINEERING
CENTER

DTIC EICE COPY

CRDEC-TR-88083

ESTIMATION OF REACTIVITY
BASED UPON STERIC FACTORS:
AMIDIZATION OF ESTERS

DTIC
ELECTE
JUN 02 1988
S D

by George R. Famini
Joseph M. Leonard, Ph.D.
RESEARCH DIRECTORATE

March 1988

"Original contains color
plates: All DTIC reproductions
will be in black and
white"

U.S. ARMY
ARMAMENT
MUNITIONS
CHEMICAL COMMAND



Aberdeen Proving Ground, Maryland 21010-5423

This document has been approved
for public release and sale; its
distribution is unlimited.

00 0000 025

Disclaimer

The findings in this report are not to be construed as an official Department of the Army position unless so designated by other authorizing documents.

Distribution Statement

Approved for public release; distribution is unlimited.

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE

REPORT DOCUMENTATION PAGE

1a. REPORT SECURITY CLASSIFICATION UNCLASSIFIED			1b. RESTRICTIVE MARKINGS	
2a. SECURITY CLASSIFICATION AUTHORITY			3. DISTRIBUTION / AVAILABILITY OF REPORT Approved for public release; distribution is unlimited.	
2b. DECLASSIFICATION / DOWNGRADING SCHEDULE			5. MONITORING ORGANIZATION REPORT NUMBER(S)	
4. PERFORMING ORGANIZATION REPORT NUMBER(S) CRDEC-TR-88083			7a. NAME OF MONITORING ORGANIZATION	
6a. NAME OF PERFORMING ORGANIZATION CRDEC		6b. OFFICE SYMBOL (If applicable) SMCCR-RSP-C		7b. ADDRESS (City, State, and ZIP Code)
6c. ADDRESS (City, State, and ZIP Code) Aberdeen Proving Ground, MD 21010-5423			9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER	
8a. NAME OF FUNDING / SPONSORING ORGANIZATION CRDEC		8b. OFFICE SYMBOL (If applicable) SMCCR-RSP-C		10. SOURCE OF FUNDING NUMBERS
8c. ADDRESS (City, State, and ZIP Code) Aberdeen Proving Ground, MD 21010-5423			PROGRAM ELEMENT NO. 14N463615	TASK NO. DE762X
11. TITLE (Include Security Classification) Estimation of Reactivity Based Upon Steric Factors: Amidization of Esters				
12. PERSONAL AUTHOR(S) Famini, George R., and Leonard, Joseph M., Ph.D.				
13a. TYPE OF REPORT Technical		13b. TIME COVERED FROM 87 May TO 87 Jul		14. DATE OF REPORT (Year, Month, Day) 1988 March
15. PAGE COUNT 16				
16. SUPPLEMENTARY NOTATION				
17. COSATI CODES			18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)	
FIELD 20	GROUP 05	SUB-GROUP	Molecular modeling, Methacrylates, Computational chemistry, polymers Molecular graphics, Esters, mjm	
19. ABSTRACT (Continue on reverse if necessary and identify by block number) Using capabilities of the Molecular Modeling Analysis and Display System (MMADS), we examined and modeled the relative reactivities of i-butyl acetate and poly i-butyl methacrylate to amidization. Due to visualization and computational constraints, the polymer has been approximated as a tetramer. The amine is sterically constrained from reaching the reactive centers of the polymer/tetramer model but freely reacts with the analogous site in the monomer. This reaction is in agreement with experimental results using both the monomer and polymer of t-butyl acetate. Keyman				
20. DISTRIBUTION / AVAILABILITY OF ABSTRACT <input checked="" type="checkbox"/> UNCLASSIFIED/UNLIMITED <input type="checkbox"/> SAME AS RPT. <input type="checkbox"/> DTIC USERS			21. ABSTRACT SECURITY CLASSIFICATION UNCLASSIFIED	
22a. NAME OF RESPONSIBLE INDIVIDUAL SANDRA J. JOHNSON			22b. TELEPHONE (Include Area Code) (301) 671-2914	22c. OFFICE SYMBOL SMCCR-SPS-T

DD FORM 1473, 84 MAR

83 APR edition may be used until exhausted.
All other editions are obsolete.SECURITY CLASSIFICATION OF THIS PAGE
UNCLASSIFIED

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE

PREFACE

The work described in this report was authorized under Project No. 14N463615DE762X. This work was started in May 1987 and completed in July 1987.

The use of trade names or manufacturers' names in this report does not constitute an official endorsement of any commercial products. This report may not be cited for purposes of advertisement.

Reproduction of this document in whole or in part is prohibited except with permission of the Commander, U.S. Army Chemical Research, Development and Engineering Center, ATTN: SMCCR-SPS-T, Aberdeen Proving Ground, Maryland 21010-5423. However, the Defense Technical Information Center and the National Technical Information Service are authorized to reproduce the document for U.S. Government purposes.

This report has been approved for release to the public.

Accession For	
NTIS GRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	



"Original contains color plates: All DTIC reproductions will be in black and white"

Blank

CONTENTS

	Page
1. INTRODUCTION	7
2. EXPERIMENTAL PROCEDURE	8
3. RESULTS	8
4. CONCLUSION	8

LIST OF FIGURES

Figure

1. Amidization Reactants	11
2. Amidization Products	12
3. Tetramer of i-Butyl Acetate (space fill)	13
4. Solvent Accessible Surface of i-Butyl Acetate Tetramer	14
5. i-Butyl Acetate (space fill)	15
6. Solvent Accessible Surface of i-Butyl Acetate	16

Blank

ESTIMATION OF REACTIVITY BASED UPON STERIC FACTORS: AMIDIZATION OF ESTERS

1. INTRODUCTION

To determine if two compounds will react based upon purely steric parameters is fairly easy to conceptualize. In principle, it should be possible to "map" the reactive site of the larger reactant such that the size of the smaller reactant is taken into effect. In this way, the steric fit of the two reactants can be examined, and the reactivity based upon steric grounds can be estimated.

The particular reaction examined was the amidization of i-butyl acetate with i-propyl amine. The reactants are shown in Figure 1, i-propyl amine on the left and i-butyl acetate on the right. Each atom is represented as a sphere with a radius equal to the corresponding van der Waals radius. In this representation, nitrogen is blue, carbon grey, oxygen red, and hydrogen is white. The end products of the reaction are displayed in Figure 2, i-propyl acetamide on the left and i-butyl alcohol (represented as the oxy anion) on the right.

One particular option within the Molecular Modeling Analysis and Display System (MMADS) at Research Directorate, U.S. Army Chemical Research, Development and Engineering Center (CRDEC), lends itself very well to this type of study. This routine, the Connolly Solvent Accessible Surface Algorithm,¹ calculates the surface of a substrate molecule in contact with a solvent of a given radius. For a reaction to take place, the i-propyl amine must reach the active site of the i-butyl acetate (a carbonyl carbon). This routine can be used to determine if it is sterically possible for a reaction to occur.

As with all modeling efforts, it is impossible (both in terms of computer time necessary and viewability) to duplicate the exact system being examined. Therefore, approximations have been made. Using an empirical optimization technique such as molecular mechanics permits large molecules (between 100 to 200 atoms) to be sterically optimized in a minimal amount of time. Using pictorial techniques, the size of the structure must be kept reasonable to permit the eye to observe the regions of interest. For these reasons, a tetramer of the i-butyl methacrylate was considered sufficient to model the polymer interactions but was small enough to allow for ready identification of the active sites.

¹Connolly, M., Science Vol. 221, p. 709 (1981).

2. EXPERIMENTAL PROCEDURE

The structures of i-butyl acetate, the corresponding tetramer, and i-propyl amine were geometrically optimized using the Allinger molecular mechanics algorithm.² The solvent accessible surface was calculated using the Connolly algorithm.¹ Both routines are resident in the MMADS, running on a VAX computer under the VMS operating system at CRDEC.³ All photographs were taken from an Adage 3000 Color Raster Display using a Dunn 635 camera.

3. RESULTS

This investigation was to determine if poly i-butyl methacrylate would undergo amidization. Figure 3 shows the optimized structure of the i-butyl acetate tetramer. The color scheme of the previous figures is used, except that the active site (the carbonyl carbon) is green for clarity. By assuming that the i-propyl amine molecule can be approximated as a sphere with a radius of 3.616 Å (the effective van der Waals radius of the molecule), the solvent accessible surface can be generated for the potential reaction pair, i-propyl amine and poly i-butyl methacrylate. This surface is shown in Figure 4. The presence of green dots, denoting that the carbonyl carbon can be "touched" by the i-propyl amine, is an indication that the reaction could occur based upon steric constraints. As seen from the figure, no green dots are visible, indicating that poly i-butyl methacrylate would be unreactive towards i-propyl amine. This agrees with experimental results.

Experimental results show, however, that the monomer unit of i-butyl acetate readily undergoes amidization. Therefore, if this model is to be successful in predicting steric based reactivity, it must also be able to predict that this reaction can occur. The monomer unit is shown in Figure 5 using the same color scheme as Figure 3. The solvent accessible surface was calculated using the same conditions as the polymer, producing the results shown in Figure 6. The significant number of green dots on the surface indicates that the carbonyl carbon in the monomer is accessible to the i-propyl amine; and therefore, a reaction can occur.

4. CONCLUSION


The study shows that it is possible to use computational techniques to rationalize apparently anomalous experimental results. In reactions where steric constraints are the primary factors, this methodology should be applicable.

²Allinger, N., and Burkert, V., ACS Monograph 177, American Chemical Society, Washington, DC (1982).

³Leonard, J., A User's Guide to the Molecular Modeling Analysis and Display System (MMADS), CRDEC-TR-86039, U.S. Army Chemical Research, Development and Engineering Center, Aberdeen Proving Ground, MD, May 1986, UNCLASSIFIED Report.

First, the solvent accessible surface algorithm assumes that the solvent molecule can be represented as a sphere. This assumption is not completely valid, and the results become less useful as the solvent molecules become less spherical. However, if the solute is much larger than the solvent, the approximation will hold. Second, this model addresses only the steric factors in a reaction, completely ignoring the electronic factors. Although this may not be important in this reaction, electronic interactions may be the rate limiting step in other reactions.

Blank






Figure 2. Amidization Products




Figure 3. Tetramer of i-Butyl Acetate (space fill)




Figure 4. Solvent Accessible Surface of i-Butyl Acetate Tetramer




Figure 5. i-Butyl Acetate (space fill)




Figure 6. Solvent Accessible Surface of i-Butyl Acetate